arXiv:1412.5748v1 [math.CV] 18 Dec 2014

THE COMPUTATION OF ZEROS OF AHLFORS MAP FOR DOUBLY CONNECTED REGIONS

ALI W.K. SANGAWI, KASHIF NAZAR, AND ALI H.M. MURID *

Abstract. The relation between the Ahlfors map and Szegö kernel $S(z,a)$ is classical. The Szegö kernel is a solution of a Fredholm integral equation of the second kind with the Kerzman-Stein kernel. The exact zeros of the Ahlfors map are unknown except for the annulus region. This paper presents a numerical method for computing the zeros of the Ahlfors map of any bounded doubly connected region. The method depends on the values of $S(z(t),a)$, $S'(z(t),a)$ and $\theta'(t)$ where $\theta(t)$ is the boundary correspondence function of Ahlfors map. A formula is derived for computing $S'(z(t),a)$. An integral equation is constructed for solving $\theta'(t)$. The numerical examples presented here prove the effectiveness of the proposed method.

1. Introduction

The conformal mapping from a multiply connected region onto the unit disk is known as the Ahlfors map. If the region is simply connected then the Ahlfors map reduces to the Riemann map. Many of the geometrical features of a Riemann mapping function are shared with Ahlfors map. The Riemann mapping function can be regarded as a solution of the following extremal problem: For a simply connected region Ω and canonical region D in the complex plane $\mathbb{C}$ and fixed a in Ω, construct an extremal analytic map

$$F : \Omega \to D \text{ with } \quad F'(a) > 0.$$

The Riemann map is the solution of this problem. It is unique conformal, one-to-one and onto map with $F(a) = 0$.

For a multiply connected region Ω of connectivity $n > 1$, the answer to the same extremal problem above becomes the Ahlfors map. It is unique analytic map

$$f : \Omega \to D$$

that is onto, $f'(a) > 0$ and $f(a) = 0$. However it has $2n-2$ branch points in the interior and is no longer one-to-one there. In fact it maps Ω onto D in an n-to-one fashion, and maps each boundary curve one-to-one onto the unit circle (see [5] and [17]). Therefore the Ahlfors map can be regarded as the Riemann mapping function in the multiply connected region.

Conformal mapping of multiply connected regions can be computed efficiently using the integral equation method. The integral equation method has been used by many authors to compute the one-to-one conformal mapping from multiply connected regions onto some standard canonical regions [3,4,6,8–11,13,15,19–23,27].

1991 *Mathematics Subject Classification.* 30C30; 30E25; 65E05.

Key words and phrases. Ahlfors map; Adjoint Neumann kernel; Generalized Neumann kernel.

*Corresponding author at: UTM Centre for Industrial and Applied Mathematics(UTM-CIAM), Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia.

Some integral equations for Ahlfors map have been given in [1,7,14,16,17]. In [3], Kerzman and Stein have derived a uniquely solvable boundary integral equation for computing the Szegö kernel of a bounded region and this method has been generalized in [1] to compute Ahlfors map of bounded multiply connected regions without relying on the zeros of Ahlfors map. In [7, 14] the integral equations for Ahlfors map of doubly connected regions requires knowledge of zeros of Ahlfors map, which are unknown in general.

In this paper, we extend the approach of Sangawi [19, 22–24] to construct an integral equation for solving $\theta'(t)$ where $\theta(t)$ is the boundary correspondence function of Ahlfors map of multiply connected region onto a unit disk.

The plan of this paper is as follows: Section 2 presents some auxiliary materials. We shall derive in Section 3, a boundary integral equation satisfied by $\theta'(t)$, where $\theta(t)$ is the boundary correspondence function of Ahlfors map of bounded multiply connected regions onto a disk. From the computed values of $\theta'(t)$, we then determine the Ahlfors map. In Section 4, we present a formula for computing the derivative of Szegö kernel which we used for finding the derivative of Ahlfors map giving another way of computing $\theta'(t)$ analytically. In Section 5, we present a method for computing the second zero of the Ahlfors map for doubly connected regions. In Section 6, we present some examples to illustrate our boundary integral equation method for finding the zeros of Ahlfors map for general doubly connected region. The numerical examples are first restricted to annulus region for which the exact Ahlfors map is known, then verified on general doubly connected region and obtained accurate results. Finally, Section 7 presents a short conclusion.

2. Auxiliary Materials

Let Ω be a bounded multiply connected region of connectivity $M+1$. The boundary Γ consists of $M+1$ smooth Jordan curves $\Gamma_0, \Gamma_1, \ldots, \Gamma_M$ such that $\Gamma_1, \ldots, \Gamma_M$ lie in the interior of Γ_0, where the outer curve Γ_0 has counterclockwise orientation and inner curves $\Gamma_1, \ldots, \Gamma_M$ have clockwise orientation. The positive direction of the contour $\Gamma = \Gamma_0 \cup \Gamma_1 \cup \cdots \cup \Gamma_M$ is usually that for which Ω is on the left as one traces the boundary as shown in Figure 1.



Figure 1. A bounded multiply connected region Ω of connectivity $M+1$.

The curves Γ_j are parameterized by 2π-periodic triple continuously differentiable complex-valued functions $z_j(t)$ with non-vanishing first derivatives

$$z_j'(t) = dz_j(t)/dt \neq 0, \quad t \in J_j = [0, 2\pi], \quad j = 0, 1, \ldots, M.$$

The total parameter domain J is defined as the disjoint union of $M+1$ intervals $J_0,\ldots,J_M$. The notation

$$z(t)=z_j(t),\quad t\in J_j,\quad j=0,1,\ldots,M.$$

must be interpreted as follows [14]: For a given $\tilde{t}\in[0,2\pi]$, to evaluate the value of $z(t)$ at $\tilde{t}$, we should know in advance the interval J_j to which $\tilde{t}$ belongs, i.e., we should know the boundary Γ_j contains $z(\tilde{t})$, then we compute $z(\tilde{t})=z_j(\tilde{t})$.

The generalized Neumann kernel formed with a complex continuously differentiable 2π-periodic function $A(t)$ for all $t\in J$, is defined by [25,26]

$$N(t,s)=\frac{1}{\pi}\mathrm{Im}\left(\frac{A(t)}{A(s)}\frac{z'(s)}{z(s)-z(t)}\right).$$

The classical Neumann kernel is the generalized Neumann kernel formed with $A(t)=1$, i.e.

$$N(t,s)=\frac{1}{\pi}\mathrm{Im}\left(\frac{z'(s)}{z(s)-z(t)}\right).$$

The kernel is continuous which takes on the diagonal the values

$$N(t,t)=\frac{1}{\pi}\left(\frac{1}{2}\mathrm{Im}\frac{z''(t)}{z'(t)}-\mathrm{Im}\frac{A'(t)}{A(t)}\right).$$

The adjoint of the function A is given by

$$\tilde{A}=\frac{z'(t)}{A(t)}.$$

The generalized Neumann kernel $\tilde{N}(s,t)$ formed with $\tilde{A}$ is given by

$$\tilde{N}(s,t)=\frac{1}{\pi}\mathrm{Im}\left(\frac{\tilde{A}(s)}{\tilde{A}(t)}\frac{z'(t)}{z(t)-z(s)}\right),$$

which implies

$$\tilde{N}(s,t)=-N^*(s,t),$$

where $N^*(s,t)=N(t,s)$ is the adjoint kernel for the generalized Neumann kernel $N(s,t)$ (see [25]).

In the remaining part of this paper, we shall only be dealing with $A(t)=1$, which the classical Neumann kernel. The adjoint kernel $N^*(s,t)$ of the classical Neumann kernel is given by

$$N^*(t,s)=N(s,t)=\frac{1}{\pi}\mathrm{Im}\left(\frac{z'(t)}{z(t)-z(s)}\right).$$

It is known that $\lambda=1$ is an eigenvalue of the kernel N with multiplicity 1 and $\lambda=-1$ is an eigenvalue of the kernel N with multiplicity M [26]. The eigenfunctions of N corresponding to the eigenvalue $\lambda=-1$ are $\left\{\chi^{[1]},\chi^{[2]},\ldots,\chi^{[M]}\right\}$, where

$$\chi^{[j]}(\xi)=\begin{cases}1, & \xi\in\Gamma_j,\\ 0, & \text{otherwise,}\end{cases}\qquad j=1,2,\ldots,M.$$

Let H^* be the space of all real Hölder continuous 2π- periodic functions $\omega(t)$ of the parameter t on J_j for $j=0,1,\ldots,M$, i.e.

$$\omega(t)=\omega_k(t),\ t\in J_j,\ k=0,1,\ldots,M,j=0,1,\ldots,M.$$

Define also the integral operator $\mathbf{J}$ by [20, 22]

$$\mathbf{J}\nu = \int_J \frac{1}{2\pi} \sum_{j=1}^{M} \chi^{[j]}(s)\chi^{[j]}(t)\nu(s)ds.$$

We also define the Fredholm integral operator N^* by

$$\mathbf{N}^*\psi(t) = \int_J N^*(t,s)\psi(s)ds, \ t \in J.$$

A complex-valued function $P(z)$ is said to satisfy the interior relationship if $P(z)$ is analytic in Ω and satisfies the non-homogeneous boundary relationship

$$P(z) = \frac{b(z)\overline{T(z)}}{\overline{G(z)}}\overline{P(z)} + \overline{H(z)}, \quad z \in \Gamma, \tag{2.1}$$

where $G(z)$ is analytic in Ω and Hölder continuous on Γ, $G(z) \neq 0$ on Γ, and $T(z) = z'(t)/|z'(t)|$ denotes the unit tangent in the direction of increasing parameters at the point $z = z(t) \in \Gamma$. The boundary relationship (2.1) also has the following equivalent form:

$$G(z) = \overline{b(z)}T(z)\frac{P(z)^2}{|P(z)|^2} + \frac{G(z)H(z)}{\overline{P(z)}}, \quad z \in \Gamma. \tag{2.2}$$

The following theorem gives an integral equation for an analytic function satisfying the non-homogeneous boundary relationship (2.1) or (2.2).

Theorem 2.1. *[19] If the function $P(z)$ satisfies the interior relationship with (2.1) or (2.2), then*

$$T(z)P(z) + \int_\Gamma K(z,w)T(w)P(w)|dw| + b(z)\left[\sum_{a_j \, inside \, \Gamma} \underset{w=a_j}{Res} \frac{P(w)}{(w-z)G(w)}\right]^- = -\overline{L_R^-(z)},$$

where

$$K(z,w) = \frac{1}{2\pi \mathrm{i}}\left[\frac{T(z)}{z-w} - \frac{b(z)\overline{T(w)}}{b(w)(\overline{z}-\overline{w})}\right],$$

and

$$L_R^-(z) = \frac{-1}{2}\frac{H(z)}{T(z)} + \mathrm{PV}\frac{1}{2\pi \mathrm{i}}\int_\Gamma \frac{\overline{b(z)}H(w)}{\overline{b(w)}(w-z)T(w)}dw.$$

The symbol " $-$ " in the superscript denotes the complex conjugate and the sum is over all those zeros that lie inside Ω. If G has no zeros in Ω, then the term containing the residue will not appear.

3. Integral equation method for computing θ'

Let $f(z)$ be the Ahlfors function which maps Ω conformally onto a unit disc. The mapping function f is determined up to a factor of modulus 1. The function f could be made unique by prescribing that

$$f(a_j) = 0, \quad f'(a_0) > 0, \quad j = 0, 1, 2, \ldots, M.$$

where $a_j \in \Omega, j = 0, 1, \ldots, M$ are the zeros of the Ahlfors map. The boundary values of f can be represented as

$$f(z_j(t)) = e^{\mathrm{i}\theta_j(t)}, \quad \Gamma_j : z = z_j(t), \quad 0 \le t \le \beta_j, \tag{3.1}$$

where $\theta_j(t), j = 0, 1, \ldots, M$ are the boundary correspondence functions of Γ_j. Also we have

$$\frac{f'(z(t))z'(t)}{f(z(t))} = \mathrm{i}\theta'(t), \tag{3.2}$$

The unit tangent to Γ at $z(t)$ is denoted by $T(z(t)) = z'(t)/|z'(t)|$. Thus it can be shown that

$$f(z_j(t)) = \frac{1}{\mathrm{i}} T(z_j(t)) \frac{\theta_j'(t)}{|\theta_j'(t)|} \frac{f'(z_j(t))}{|f'(z_j(t))|}, \qquad z_j \in \Gamma_j. \tag{3.3}$$

By the angle preserving property of conformal map, the image of Γ_0 remains in counter-clockwise orientation so $\theta_0'(t) > 0$ while the images of inner boundaries Γ_j in clockwise orientation so $\theta_j'(t) < 0$, for $j = 1, \ldots, M$. Thus

$$\operatorname{sign}(\theta_j'(t)) = \begin{cases} +1, & j = 0, \\ -1, & j = 1, \ldots, M. \end{cases}$$

The boundary relationship (3.3) can be written briefly as

$$f(z) = \operatorname{sign}(\theta'(t)) \frac{1}{\mathrm{i}} T(z) \frac{f'(z)}{|f'(z)|}, \quad z \in \Gamma. \tag{3.4}$$

Since the Ahlfors map can be written as $f(z) = \prod_{j=0}^{M} (z - a_j)\hat{g}(z)$, where $\hat{g}(z)$ is analytic in Ω and $\hat{g}(z) \neq 0$ in Ω, we have

$$\frac{f'(z)}{f(z)} = \frac{\hat{g}'(z)}{\hat{g}(z)} + \sum_{j=0}^{M} \frac{1}{z - a_j}, \quad z \in \Gamma, \tag{3.5}$$

so that

$$D(z) = \frac{f'(z)}{f(z)} - \sum_{j=0}^{M} \frac{1}{z - a_j} = \frac{\hat{g}'(z)}{\hat{g}(z)}. \tag{3.6}$$

is analytic in Ω. Squaring both sides of (3.4), gives

$$f(z)^2 = -T(z)^2 \frac{f'(z)^2}{|f'(z)|^2}, \tag{3.7}$$

which implies

$$\frac{f'(z)}{f(z)} = -\overline{T(z)}^2 \overline{\left(\frac{f'(z)}{f(z)} \right)}. \tag{3.8}$$

From (3.6), we have

$$\frac{f'(z)}{f(z)} = D(z) + \sum_{j=0}^{M} \frac{1}{z - a_j}. \tag{3.9}$$

Substituting (3.9) into (3.8)

$$D(z) + \sum_{j=0}^{M} \frac{1}{z - a_j} = -\overline{T(z)}^2 \overline{D(z)} - \sum_{j=0}^{M} \frac{\overline{T(z)}^2}{\overline{z} - \overline{a_j}} \tag{3.10}$$

or

$$D(z) = -\overline{T(z(t))}^2 \overline{D(z)} - \sum_{j=0}^{M} \frac{\overline{T(z)^2}}{\overline{z} - \overline{a_j}} - \sum_{j=0}^{M} \frac{1}{z - a_j}. \tag{3.11}$$

Comparing (3.11) with (2.1), we get

$$P(z) = D(z), b(z) = -\overline{T(z)}, G(z) = 1, H(z) = -\sum_{j=0}^{M} \frac{T(z)^2}{z - a_j} - \sum_{j=0}^{M} \frac{1}{\overline{z} - \overline{a_j}}. \tag{3.12}$$

Substituting these assignments into Theorem 2.1, we obtain

$$T(z)D(z) + \int_{\Gamma} K(z, w)T(w)D(w)|dw| + 0 = -\overline{L_R^-(z)}, \tag{3.13}$$

where

$$K(z, w) \quad = \frac{1}{2\pi \mathrm{i}} \left[\frac{T(z)}{z - w} - \frac{\overline{T(z)}}{\overline{z} - \overline{w}} \right] \quad = \frac{1}{\pi} \mathrm{Im} \left(\frac{T(z)}{z - w} \right) = N(z, w)$$

and

$$L_R^-(z) = \frac{-1}{2T(z)} \left[\sum_{j=0}^{M} \frac{T(z)^2}{z - a_j} - \sum_{j=0}^{M} \frac{1}{\overline{z} - \overline{a_j}} \right] + \mathrm{PV} \frac{1}{2\pi \mathrm{i}} \int_{\Gamma} \frac{T(z)}{(w - z)T(w)^2} \left[\sum_{j=0}^{M} \frac{T(w)^2}{w - a_j} - \sum_{j=0}^{M} \frac{1}{\overline{w} - \overline{a_j}} \right].$$

Using the fact that [20]

$$\mathrm{PV} \frac{1}{2\pi \mathrm{i}} \int_{\Gamma} \frac{1}{(w - z)(w - a_j)} dw = -\frac{1}{2(z - a_j)},$$

$dw = T(w)|dw|$, and after some simplifications, (3.13) becomes

$$T(z) \frac{f'(z)}{f(z)} + \int_{\Gamma} N(z, w) \frac{f'(w)}{f(w)} T(w)|dw| = 2\mathrm{i}\mathrm{Im} \left[\sum_{j=0}^{M} \frac{T(z)}{z(t) - a_j} \right], \tag{3.14}$$

where

$$N(z,w)=\begin{cases}\dfrac{1}{2\pi \mathrm{i}}\left[\dfrac{T(z)}{z-w}-\dfrac{\overline{T(z)}}{\overline{z}-\overline{w}}\right], & z\neq w\in\Gamma,\\ \dfrac{1}{2\pi|z'(t)|}\mathrm{Im}\left(\dfrac{z''(t)}{z'(t)}\right), & z=w\in\Gamma,\end{cases} \tag{3.15}$$

with

$$\frac{1}{2\pi \mathrm{i}}\int_{-\Gamma_j}\frac{f'(w)T(w)}{f(w)}|dw|=1,\quad j=1,2,\ldots,M. \tag{3.16}$$

In integral equation (3.14), letting $z=z(t)$, $w=z(s)$ and multiplying both sides by $|z'(t)|$, gives

$$\frac{f'(z(t))}{f(z(t))}z'(t)+\int_{\Gamma}N(z(t),z(s))\frac{f'(z(s))}{f(z(s))}z'(s)ds=2\mathrm{iIm}\left[\sum_{j=0}^{M}\frac{z'(t)}{z(t)-a_j}\right]. \tag{3.17}$$

Using (3.2), the above equation becomes

$$\theta'(t)+\int_{J}N(t,s)\theta'(s)ds=2\mathrm{Im}\left[\sum_{j=0}^{M}\frac{z'(t)}{z(t)-a_j}\right]. \tag{3.18}$$

Since $N^*(s,t)=N(t,s)$, the integral equation (3.18) in the operator form is

$$(\mathbf{I}+\mathbf{N}^*)\theta'_j=\phi,\qquad j=0,1,\ldots,M, \tag{3.19}$$

where

$$\phi=2\mathrm{Im}\left[\sum_{j=0}^{M}\frac{z'(t)}{z(t)-a_j}\right]. \tag{3.20}$$

By Theorem 12 in [26], $\lambda=-1$ is an eigenvalue of N^* with multiplicity M, therefore the integral equation (3.19) is not solvable. To overcome this problem, we note from (3.16) and (3.2) that

$$\mathbf{J}\theta'=\left(0,\frac{1}{2\pi}\int_{J_1}\theta'(s)ds,\frac{1}{2\pi}\int_{J_2}\theta'(s)ds,...,\frac{1}{2\pi}\int_{J_M}\theta'(s)ds\right)$$

which implies that

$$\mathbf{J}\theta'=\psi, \tag{3.21}$$

where

$$\psi=(0,1,1,\ldots,1). \tag{3.22}$$

By adding (3.19) and (3.21), we get

$$(\mathbf{I}+\mathbf{N}^*+\mathbf{J})\theta'=\phi+\psi \tag{3.23}$$

which implies

$$\phi=(\mathbf{I}+\mathbf{N}^*+\mathbf{J})\theta'-\psi. \tag{3.24}$$

In the next section, we derive another formula for computing θ' from which the second zero of the Ahlfors map for doubly connected region can be calculated using (3.24).

4. An Alternative Formula for Computing θ'

The Ahlfors map is related to the Szegö kernel $S(z,a_0)$ and the Garabedian kernel $L(z,a_0)$ by [1]

$$f(z)=\frac{S(z,a_0)}{L(z,a_0)},\quad z\in\Omega\cup\Gamma. \tag{4.1}$$

The Szegö kernel $S(z,a_0)$ and Garabedian kernel $L(z,a_0)$ are related on Γ as

$$L(z,a_0)=-\mathrm{i}\overline{T(z)S(z,a_0)},\quad z\in\Gamma,$$

so that (4.1) becomes

$$f(z)=\frac{1}{\mathrm{i}}\frac{S(z,a_0)T(z)}{\overline{S(z,a_0)}},\quad z\in\Gamma, \tag{4.2}$$

or

$$f(z(t))=\frac{1}{\mathrm{i}}\frac{S(z(t),a_0)T(z(t))}{\overline{S(z(t),a_0)}},\quad z(t)\in\Gamma. \tag{4.3}$$

From Bell [1], we have the integral equation for the Szegö kernel

$$S(z,a_0)+\int_\Gamma A(z,w)S(w,a_0)|dw|=g(z),\quad z\in\Gamma, \tag{4.4}$$

where

$$A(z,w)=\begin{cases}\dfrac{1}{2\pi\mathrm{i}}\left[\dfrac{T(w)}{z-w}-\dfrac{\overline{T(z)}}{\overline{z-w}}\right], & z\neq w\in\Gamma\\ 0, & z=w,\end{cases}$$

and

$$g(z(t))=-\frac{1}{2\pi\mathrm{i}}\frac{\overline{T(z(t))}}{\overline{z(t)-a_0}}. \tag{4.5}$$

With $z=z(t)$ and $w=z(s)$, (4.4) becomes

$$S(z(t),a_0)+\int_{J_j}A(z(t),z(s))S(z(s),a_0)|z'(s)|ds=g(z(t)).$$

Differentiate both sides with respect to t, we get

$$\frac{d}{dt}S(z(t),a_0)+\frac{d}{dt}\int_{J_j}A(z(t),z(s))S(z(s),a_0)|z'(s)|ds=\frac{d}{dt}g(z(t)),$$

which is equivalent to

$$S'(z(t),a_0)z'(t)=g'(z(t))z'(t)-\int_{J_j}\left[\frac{d}{dt}A(z(t),z(s))z'(t)\right]S(z(s),a_0)|z'(s)|ds. \tag{4.6}$$

Differentiate (4.5) with respect to t, gives

$$g'(z(t))z'(t)=-\frac{1}{2\pi\mathrm{i}}\left[\frac{\overline{T'(z(t))z'(t)}}{\overline{z(t)-a_0}}-\frac{\overline{z'(t)}\ \overline{T(z(t))}}{\overline{(z(t)-a_0)^2}}\right]. \tag{4.7}$$

Since

$$T(z(t))=\frac{z'(t)}{|z'(t)|},$$

we get

$$T'(z(t))z'(t) = \frac{z''(t)}{2|z'(t)|} - \frac{z'^2(t)\overline{z''(t)}}{2|z'(t)|^3}. \tag{4.8}$$

Since

$$A(z(t), z(s)) = \frac{1}{2\pi \mathrm{i}} \left[\frac{T(z(s))}{z(t) - z(s)} - \frac{\overline{T(z(t))}}{\overline{z(t) - z(s)}} \right], \quad \text{for } t \neq s,$$

differentiating with respect to t, we get

$$\frac{d}{dt} A(z(t), z(s)) = \frac{1}{2\pi \mathrm{i}} \left[\frac{-z'(t)\ T(z(s))}{(z(t) - z(s))^2} - \frac{\overline{T'(z(t))z'(t)}}{(\overline{z(t)} - \overline{z(s)})} + \frac{\overline{T(z(t))}\ \overline{z'(t)}}{(\overline{z(t)} - \overline{z(s)})^2} \right],$$

which implies

$$\begin{aligned} 2\pi \mathrm{i}|z'(s)| \frac{d}{dt} A(z(t), z(s)) = & - \left(\frac{z'(t)}{z(t) - z(s)} \right)^2 \left(\frac{z'(s)}{z'(t)} \right) \\ & - \left(\frac{\overline{z''(t)}z'(t) - \overline{z'(t)}z''(t)}{2|z'(t)|^2} \right) \left(\overline{\frac{z'(t)}{z(t) - z(s)}} \right) \left| \frac{z'(s)}{z'(t)} \right| \\ & + \left(\overline{\frac{z'(t)}{z(t) - z(s)}} \right)^2 \left| \frac{z'(s)}{z'(t)} \right|. \end{aligned} \tag{4.9}$$

For s sufficiently close to t, we have

$$z(s) = z(t) + z'(t)(s-t) + \frac{1}{2} z''(t)(s-t)^2 + \frac{1}{6} z'''(t)(s-t)^3 + O((s-t)^4)$$

which implies

$$\frac{z'(t)}{z(s) - z(t)} = \frac{1}{s-t} \left[1 + \frac{1}{2} \frac{z''(t)}{z'(t)} (s-t) + \frac{1}{6} \frac{z'''(t)}{z'(t)} (s-t)^2 + O((s-t)^4) \right]^{-1} \tag{4.10}$$

Using the fact that

$$\frac{1}{1+\varrho} = 1 - \varrho + \varrho^2 + O(\varrho^3), \tag{4.11}$$

for ϱ close to zero, (4.10) becomes

$$\frac{z'(t)}{z(s) - z(t)} = \frac{1}{s-t} - \frac{1}{2} \frac{z''(t)}{z'(t)} - \frac{1}{6} \frac{z'''(t)}{z'(t)} (s-t) + \frac{1}{4} \frac{z''^2(t)}{z'^2(t)} (s-t) + O((s-t)^2). \tag{4.12}$$

We observe that

$$\frac{z'(s)}{z'(t)} = 1 + \frac{z''(t)}{z'(t)} (s-t) + \frac{1}{2} \frac{z'''(t)}{z'(t)} (s-t)^2 + O((s-t)^3) \tag{4.13}$$

Next observe that

$$\begin{aligned} \left| \frac{z'(s)}{z'(t)} \right|^2 = 1 & + 2\mathrm{Re} \left(\frac{z''(t)}{z'(t)} \right) (s-t) \\ & + \left(\mathrm{Re} \left(\frac{z'''(t)}{z'(t)} \right) + \left| \frac{z''(t)}{z'(t)} \right|^2 \right) (s-t)^2 + O((s-t)^3) \end{aligned} \tag{4.14}$$

Using the fact that

$$\sqrt{1 + \hat{x}} = 1 + \frac{1}{2}\hat{x} - \frac{1}{8}\hat{x}^2 + \cdots \tag{4.15}$$

for small $\hat{x}$, (4.14) becomes

(4.16)
$$\left|\frac{z'(s)}{z'(t)}\right| = 1+\mathrm{Re}\left(\frac{z''(t)}{z'(t)}\right)(s-t)+\frac{1}{2}\left(\mathrm{Re}\left(\frac{z'''(t)}{z'(t)}\right)+\left|\frac{z''(t)}{z'(t)}\right|^2\right)(s-t)^2+O((s-t)^3).$$

Substituting (4.16), (4.13), (4.12) into (4.9) and then taking the limit $s \to t$ to both sides, we obtain

$$\frac{d}{dt}A(z(t),z(s)) = \begin{cases} \dfrac{1}{2\pi \mathrm{i}}\left[\dfrac{-z'(t)\ T(z(s))}{(z(t)-z(s))^2} - \dfrac{\overline{T'(z(t))}\ \overline{z'(t)}}{(\overline{z(t)}-\overline{z(s)})} + \dfrac{\overline{T(z(t))}\ \overline{z'(t)}}{(\overline{z(t)}-\overline{z(s)})^2}\right], & t \neq s \in \Gamma, \\ \dfrac{1}{12\pi\,|z'(t)|}\mathrm{Im}\left(\dfrac{z'''(t)}{z'(t)}\right), & t = s \in \Gamma. \end{cases}$$

Using the values from (4.4) and (4.7), we can find the derivative of Szegö kernel from (4.6). By defining the following terms,

$$\begin{aligned} &f_p = f'(z(t))z'(t), \quad S_p = S'(z(t),a_0)z'(t), \\ &S_z = S(z(t),a_0), \quad T_p = T'(z(t))z'(t), \quad T_z = T(z(t)). \end{aligned} \tag{4.17}$$

and differentiating both sides of (4.3) with respect to t, the result is

$$f_p = \frac{1}{\mathrm{i}}\left[\frac{S_p\ T_z + S_z\ T_p}{\overline{S_z}} - \frac{\overline{S_p}\ (S_z T_z)}{\overline{(S_z)^2}}\right]. \tag{4.18}$$

By using the values from (4.2) and (4.18) in (3.2), we get

$$\theta'(t) = 2\mathrm{Im}\left(\frac{S_p}{S_z}\right) + \mathrm{Im}\left(\frac{z''(t)}{z(t)}\right) \tag{4.19}$$

which is the second formula for computing θ'.

5. Computing the Second Zero of the Ahlfors Map for Doubly Connected Region

In particular, if Ω is a doubly connected region, i.e. $M = 1$, then (3.20) becomes

$$\phi = 2\mathrm{Im}\left[\frac{z'(t)}{z(t)-a_0} + \frac{z'(t)}{z(t)-a_1}\right]. \tag{5.1}$$

As ϕ is known from (3.24), and the zero $a_0 \in \Omega$ can be freely prescribed, the only unknown in (5.1) is the second zero a_1 of Ahlfors map. The above equation can be written as

$$\mathrm{Im}\left[\frac{z'(t)}{z(t)-a_1}\right] = k_1(t). \tag{5.2}$$

where

$$k_1(t) = \frac{1}{2}\left[\phi - 2\mathrm{Im}\left(\frac{z'(t)}{z(t)-a_0}\right)\right]. \tag{5.3}$$

Now we suppose that

$$\begin{aligned} z(t) &= x(t) + \mathrm{i}y(t), \\ a_1 &= \alpha + \mathrm{i}\beta. \end{aligned}$$

Then (5.2) becomes

$$\frac{y'(t)(x(t)-\alpha)+x'(t)(\beta-y(t))}{(x(t)-\alpha)^2+(y(t)-\beta)^2}=k_1(t) \tag{5.4}$$

After some algebraic manipulations, we obtain

$$k_2(t)\alpha+k_3(t)\beta+k_1(t)(\alpha^2+\beta^2)=-k_4(t), \tag{5.5}$$

where

$$k_2(t) = y'(t)-2k_1(t)x(t), \tag{5.6}$$

$$k_3(t) = -2k_1(t)y(t)-x'(t), \tag{5.7}$$

$$k_4(t) = k_1(t)x^2+k_1(t)y^2+x'(t)y(t)-y'(t)x(t). \tag{5.8}$$

Equation (5.5) can be viewed as containing three unknowns namely α, β and $\alpha^2+\beta^2$. To determine them, we need three equations. For this we can choose any three values of parameter t in the given interval. At these three random values of t, we can find the values of the coefficients $k_1(t)$, $k_2(t)$, $k_3(t)$ and $k_4(t)$. By solving the obtained system of three equations with three unknowns, we can find the value of α and β, which are the real and imaginary parts of the second zero of Ahlfors map for doubly connected region. By solving the obtained system of three equations with three unknowns, we can find the value of $a_1=\alpha+\mathrm{i}\beta$, which is the second zero of Ahlfors map for doubly connected region.

6. Numerical Examples

For solving the integral equation (4.4) numerically, the reliable procedure is by using the Nyström method with the trapezoidal rule with n equidistant nodes in each interval J_j, $j=1,\ldots,M$ [8–11, 13]. The trapezoidal rule is the most accurate method for integrating periodic functions numerically [2, pp.134-142]. By solving the integral equation (4.4) for $S(z(t),a_0)$ gives $S_p=S'(z(t),a_0)z'(t)$ from (4.6) and $\theta'(t)$ from (4.19). By (3.24) we get the value of ϕ and then from (5.1) we get the value of $a_1=\alpha+\mathrm{i}\beta$. We then apply (4.3) and the Cauchy integral formula to compute $f(\alpha+\mathrm{i}\beta)$. For evaluating the Cauchy integral formula $f(z)=(1/(2\pi\mathrm{i}))\int_\Gamma f(w)/(w-z)dw$ numerically, we use the equivalent form

$$f(z)=\frac{\int_\Gamma \frac{f(w)}{w-z}dw}{\int_\Gamma \frac{1}{w-z}dw}, \quad z\in\Omega, \tag{6.1}$$

which also works very well for $z\in\Omega$ near the boundary Γ. When the trapezoidal rule is applied to the integrals in (6.1), the term in the denominator compensates the error in the numerator (see [22]). All the computations were done using MATLAB 7.12.0.635(R2011a).

Example 6.1. Consider an annulus region bounded by the two circles

$$\Gamma_0:\{z(t) = e^{\mathrm{i}t}\},$$
$$\Gamma_1:\{z(t) = re^{-\mathrm{i}t}\}, \qquad t:0\le t\le 2\pi, 0<r<1.$$

In [17], Tegtmeyer and Thomas computed the Ahlfors map using Szegö and Garabedian kernels for the annulus region, where the authors have used series representations of both Szegö kernel and Garabedian kernel. With these representations,

they found the two zeros a_0 and $a_1 = \dfrac{-r}{\overline{a_0}}$ for Ahlfors map, where r is the radius of the inner circle. They have also considered the symmetry case where the zeros are $a_0 = \sqrt{r}$ and $a_1 = -\sqrt{r}$. This example has also been considered in [14] where Ahlfors map was computed using a boundary integral equation related to a Riemann-Hilbert problem. Here we shall use these values of zeros of Ahlfors map for comparison with our numerical zeros with proposed method in the annulus $r < |z| < 1$. In this example we have chosen $r = 0.1$ and the first zero $a_0 = 0.6$. See Table 1 for numerical comparison of the computed second zeros a_{1n} from (5.5) and the exact second zeros of the Ahlfors map, $a_1 = -r/\overline{a_0}$.

TABLE 1. Maximum error norm $||a_{1n} - a_1||$ with $r = 0.1$ for Example 6.1.

Non-Symmetric case $(a_0 = 0.5, a_1 = -r/\overline{a_0})$	
n	$\|\|a_{1n} - a_1\|\|$
64	2.66(-15)
Symmetric case $(a_0 = \sqrt{r}, a_1 = -\sqrt{r})$	
n	$\|\|a_{1n} - a_1\|\|$
64	5.56(-14)

Example 6.2. Consider a doubly connected region Ω bounded by the two non-concentric circles

$$\begin{aligned} \Gamma_0 : \{z(t) &= 2e^{\mathrm{i}t}\}, \\ \Gamma_1 : \{z(t) &= c + re^{-\mathrm{i}t}\}, \qquad t : 0 \le t \le 2\pi, \end{aligned}$$

with $c = 0.2+0.6\mathrm{i}$ and radius $r = 0.3$. The test region is shown in Figure 2. Given a first zero a_0 of the Ahlfors map, the exact second zero a_1 is unknown for this region. We compute a_{1n} from (5.5) which is the approximate value of a_1. The acuracy is measured by computing $f(a_{1n})$ from (6.1). The theoretical value of $f(a_1)$ is zero. See Table 2 for the numerical values of a_{1n} and $f(a_{1n})$ involving two choices of a_0.



FIGURE 2. The region Ω for Example 6.2.

TABLE 2. Numerical values of a_{1n} and $f(a_{1n})$ for Example 6.2.

$a_0 = -0.5 + 0.2\mathrm{i}$		
n	a_{1n}	$f(a_{1n})$
16	$0.7135 + 1.0357\mathrm{i}$	$2.2(-03)$
32	$0.7125 + 1.0342\mathrm{i}$	$1.12(-06)$
64	$0.7125 + 1.0342\mathrm{i}$	$2.9(-13)$
128	$0.7125 + 1.0342\mathrm{i}$	$4.8(-15)$
$a_0 = -\mathrm{i}$		
n	a_{1n}	$f(a_{1n})$
16	$0.2831 + 1.0284\mathrm{i}$	$1.0(-01)$
32	$0.2763 + 1.0031\mathrm{i}$	$7.3(-04)$
64	$0.2763 + 1.0031\mathrm{i}$	$2.7(-08)$
128	$0.2763 + 1.0031\mathrm{i}$	$4.6(-15)$

Example 6.3. Consider a doubly connected region Ω bounded by Γ_0 and Γ_1

$$\begin{aligned}\Gamma_0 : \{z(t) &= -0.1 - 0.4i + (6 + 0.8\cos(18t))e^{(\mathrm{i}t)}\},\\ \Gamma_1 : \{z(t) &= \xi + (1.2 + 0.4\cos(4t))e^{(-\mathrm{i}t)}\}, \qquad t : 0 \le t \le 2\pi,\end{aligned}$$

The test regions are shown in Figure 3 for two different values of ξ. Given a first zero a_0 of the Ahlfors map, the exact second zero a_1 is also unknown for this region. We compute a_{1n} from (5.5) which is the approximate value of a_1. The acuracy is measured by computing $f(a_{1n})$ from (6.1). The theoretical value of $f(a_1)$ is zero. See Table 3 for the numerical values of a_{1n} and $f(a_{1n})$.



(a) Ω ($\xi = 0.6452 - 0.8655\mathrm{i}$) (b) Ω ($\xi = -2.4516 + 2.3626\mathrm{i}$)

FIGURE 3. The region Ω for Example 6.3.

TABLE 3. Numerical values of a_{1n} and $f(a_{1n})$ for Example 6.3.

$\xi = 0.6452 - 0.8655\text{i}$ & $a_0 = -1.3088 + 1.8012\text{i}$		
n	a_{1n}	$f(a_{1n})$
128	$2.0567 - 2.4889\text{i}$	$3.0(-02)$
256	$2.0625 - 2.5444\text{i}$	$5.8(-05)$
512	$2.0624 - 2.5445\text{i}$	$8.9(-08)$
1024	$2.0624 - 2.5445\text{i}$	$9.8(-12)$
$\xi = -2.4516 + 2.3626\text{i}$ & $a_0 = 2.2673 - 2.0351\text{i}$		
n	a_{1n}	$f(a_{1n})$
256	$-3.0647 + 3.2599\text{i}$	$6.9(-04)$
512	$-3.0648 + 3.2600\text{i}$	$1.8(-07)$
1024	$-3.0648 + 3.2600\text{i}$	$5.6(-12)$

7. Conclusion

In this paper, we have constructed a numerical method for finding second zero of the Ahlfors map of doubly connected regions. We derived two formulas for the derivative of the boundary correspondence function $\theta(t)$ of the Ahlfors map. These formulas were then used to find the second zero of the Ahlfors map for any smooth doubly connected regions. Analytical method for computing the exact zeros of Ahlfors map for annulus region is presented in [16] and [17] but the problem of finding zeros for arbitrary doubly connected regions is the first time presented in this paper. The numerical examples presented have illustrated that our method involving boundary integral equation has high accuracy.

Acknowledgment

This work was supported in part by the Malaysian Ministry of Education (MOE) through the Research Management Centre (RMC), Universiti Teknologi Malaysia (FRGS Ref. No. PY/2014/04077). This support is gratefully acknowledged.

References

[1] Bell, S.R.: Numerical computation of the Ahlfors map of a multiply connected planar domain, J. Math. Anal. Appl. **120**(1), 211–217 (1986).
[2] Davis, P.J., Rabinowitz, P.: Methods of Numerical Integration, 2nd Edition, Academic Press, Orlando (1984).
[3] Kerzman N., Stein, E.M.: The Cauchy kernel, the Szegö kernel, and the Riemann mapping function, Math. Ann, **236**(1), 85–93 (1978).
[4] Kerzman N., Trummer, M.R.: Numerical conformal mapping via the Szegö kernel, J. Comput. Appl. Math, **14**(1), 111–123 (1986).
[5] Krantz, S. G.: Geometric Function Theory: Explorations in Complex Analysis, Birkhäuser, Boston (2006).
[6] Lee B., Trummer, M. R.: Multigrid conformal mapping via the Szegö kernel, Electron. Trans. Numer. Anal, **2**, 22–43 (1994).
[7] Murid, A. H. M., Razali, M. R. M.: An integral equation method for conformal mapping of doubly connected regions, Matematika, **15**(2), 79–93 (1999).
[8] Nasser, M. M. S.: A boundary integral equation for conformal mapping of bounded multiply connected regions, Comput. Methods Funct. Theory, **9**(1), 127–143(2009).
[9] Nasser, M. M. S.: Numerical conformal mapping via a boundary integral equation with the generalized Neumann kernel, SIAM J. Sci. Comput, bf31(3), 1695–1715(2009).

[10] Nasser, M. M. S.: Numerical conformal mapping of multiply connected regions onto the second, third and fourth categories of Koebe's canonical slit domains, J. Math. Anal. Appl, **382**(1), 47–56 (2011).
[11] Nasser, M. M. S.: Numerical conformal mapping of multiply connected regions onto the fifth category of Koebe's canonical slit regions, J. Math. Anal. Appl, **398**(2), 729–743 (2012).
[12] Nasser, M. M. S., Murid, A. H. M., Ismail, M., Alejaily, E.M.A.: Boundary integral equations with the generalized Neumann kernel for Laplaces equation in multiply connected regions, Applied Mathematics and Computation, **217**(9), 4710-4727(2012).
[13] Nasser, M. M. S., Al-Shihri, F. A. A.: A fast boundary integral equation method for conformal mapping of multiply connected regions, SIAM J. Sci. Comput, **35**(3), A1736–A1760 (2013).
[14] Nasser, M. M. S., and Murid, A. H. M.: A boundary Integral equation with the generalised Neumann Kernel for the Ahlfors map, arXiv preprint, arXiv:1310.2260 (2013).
[15] O'Donnell, S. T., Rokhlin, V. : A fast algorithm for the numerical evaluation of conformal mappings, SIAM J. Sci. Stat. Comput, **10**(3), 475–487 (1989).
[16] Tegtmeyer, T. J.: The Ahlfors Map and Szegö Kernel in Multiply Connected Domains, PhD thesis, Purdue University (1998).
[17] Tegtmeyer, T. J., Thomas, A.D.: The Ahlfors map and Sezegö kernel for an annulus, Rocky Mountain J. Math, **29**(2), 709–723 (1999).
[18] Saff, E. B., Snider, A. D.: Fundamentals of Complex Analysis, Pearson Education Inc., New Jersey (2003).
[19] Sangawi, A. W. K.: Boundary integral equations for conformal mappings of bounded multiply connected regions. PhD thesis, Universiti Teknologi Malaysia (2012).
[20] Sangawi, A. W. K., Murid, A. H. M., Nasser, M. M. S.: Linear integral equations for conformal mapping of bounded multiply connected regions onto a disk with circular slits. Applied Mathematics and Computation, **218**(5), 2055-2068 (2011).
[21] Sangawi, A. W. K., Murid, A. H. M., Nasser, M. M. S.: Circular slits map of bounded multiply connected regions, Abstract and Applied Analysis. Hindawi Publishing Corporation, **2012**, Hindawi Publishing Corporation (2012).
[22] Sangawi, A. W. K., Murid, A. H. M.: Annulus with Spiral Slits Map and its Inverse of Bounded Multiply Connected Regions, International Journal of Science & Engineering Research, **4**(10), 1447-1454 (2013).
[23] Sangawi, A. W. K.: Spiral slits map and its inverse of bounded multiply connected regions, Applied Mathematics and Computation, **228**, 520-530 (2014).
[24] Sangawi, A. W. K.: Straight slits map and its inverse of bounded multiply connected regions, Advances in Computational Mathematics, 1–17 (2014). doi 10.1007/s10444-014-9368-x.
[25] Wegmann, R., Murid, A. H. M., Nasser, M. M. S.: The Riemann-Hilbert problem and the generalized Neumann kernel, Journal of Computational and Applied Mathematics, **182**(2), 388-415 (2005).
[26] Wegmann, R., Nasser, M. M. S.: The Riemann-Hilbert problem and the generalized Neumann kernel on multiply connected regions, J. Comput. Appl. Math, **214**(1), 36–57 (2008).
[27] Yunus, A. A. M., Murid, A. H. M., Nasser M. M. S.: Conformal mapping of unbounded multiply connected regions onto canonical slit regions, Abstract and Applied Analysis, **2012**, Hindawi Publishing Corporation (2012).

UTM Centre for Industrial and Applied Mathematics(UTM-CIAM), Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia.

Department of Mathematics, School of Science, Faculty of Science and Science Education, University of Sulaimani, 46001 Sulaimani, Kurdistan, Iraq.
E-mail address: alisangawi2000@yahoo.com (Ali W. K. Sangawi)

Department of Mathematical Sciences, Faculty of Science, Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia.

Department of Mathematics, COMSATS Institute of Information Technology, P.O.Box 54000 Defence Road Off Raiwind Road Lahore Pakistan.
E-mail address: knazar@ciitlahore.edu.pk (Kashif Nazar)

UTM Centre for Industrial and Applied Mathematics(UTM-CIAM), Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia.

Department of Mathematical Sciences, Faculty of Science, Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia.
E-mail address: alihassan@utm.my (Ali H. M. Murid)